

**ALKA-SELTZER PLUS COLD DAY AND NIGHT- aspirin, dextromethorphan hydrobromide, phenylephrine bitartrate, doxylamine succinate**  
**Bayer HealthCare LLC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Alka-Seltzer Plus® Cold Day & Night Effervescent Tablets**

**Alka-Seltzer Plus® Cold Day Effervescent Tablets**

***Drug Facts***

***Active ingredients (in each tablet)***

Aspirin 325 mg (NSAID)\*

Dextromethorphan hydrobromide 10 mg

Phenylephrine bitartrate 7.8 mg

\*nonsteroidal anti-inflammatory drug

***Purposes***

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

***Uses***

•temporarily relieves these symptoms due to a cold with cough:

□ • minor aches and pains □ • headache □ • cough

□ • nasal and sinus congestion □ • sore throat

□ • temporarily reduces fever

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

□ • hives □ • facial swelling □ • asthma (wheezing) □ • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs

(aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma • thyroid disease • diabetes
  - cough with excessive phlegm (mucus)
  - difficulty in urination due to enlargement of the prostate gland
  - persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for
  - • gout □ • diabetes □ • arthritis

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stools

- have stomach pain that does not get better
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- • fever gets worse or lasts more than 3 days
- • redness or swelling is present
- • new symptoms occur
- • ringing in the ears or loss of hearing occurs
- • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- • nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

### ***Directions***

- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
- children under 12 years: do not use

## **Other information**

### ***Other information***

- **each tablet contains:** sodium 416 mg
- Phenylketonurics: Contains Phenylalanine 9 mg Per Tablet
- store at room temperature. Avoid excessive heat

***Inactive ingredients*** acesulfame potassium, anhydrous citric acid, aspartame, calcium silicate, dimethicone, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

**Questions or comments? 1-800-986-0369** (Mon – Fri 9AM – 5PM EST)

**Alka-Seltzer Plus® Cold Night Effervescent Tablets**

### ***SPL***

### ***Active ingredients (in each tablet)***

Aspirin 500 mg (NSAID)\*

Dextromethorphan hydrobromide 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine bitartrate 7.8 mg

\*nonsteroidal anti-inflammatory drug

### ***Purposes***

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

### **Uses**

#### ***Uses***

- • temporarily relieves these symptoms due to a cold:
- • minor aches and pains □ • headache □ • runny nose
- • sinus congestion and pressure □ • cough □ • sneezing
- • sore throat □ • nasal congestion
- • temporarily reduces fever

### ***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives □ • facial swelling □ • asthma (wheezing) □ • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma
  - diabetes
  - thyroid disease
  - glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for
  - • gout □ • diabetes □ • arthritis
  - • taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- may cause marked drowsiness
  - avoid alcoholic drinks
  - alcohol, sedatives, and tranquilizers may increase drowsiness
  - be careful when driving a motor vehicle or operating machinery
  - excitability may occur, especially in children

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stools
- have stomach pain that does not get better
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## **Directions**

### ***Directions***

- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water at bedtime (may be taken every 4 to 6 hours). Do not exceed 8 tablets in 24 hours or as directed by a doctor.
- children under 12 years: do not use

## **Other information**

### ***Other information***

- each tablet contains: sodium 476 mg
- Phenylketonurics: Contains Phenylalanine 5.6 mg Per Tablet
- store at room temperature. Avoid excessive heat.

***Inactive ingredients*** acesulfame potassium, anhydrous citric acid, aspartame, calcium silicate, dimethicone, docusate sodium, flavors (natural & artificial), mannitol, povidone, sodium benzoate, sodium bicarbonate

## Questions or comments?

**Questions or comments?** 1-800-986-0369 (Mon-Fri 9AM – 5PM EST)

## Package Display Label

Alka-Seltzer

PLUS®

DAY NON-DROWSY

**ASPIRIN (NSAID)** / Pain Reliever-Fever Reducer

Dextromethorphan HBr / Cough Suppressant

Phenylephrine Bitartrate / Nasal Decongestant

- Nasal Congestion
- Headache & Body Ache
- Cough
- Sore Throat
- Sinus Pressure

12 EFFERVESCENT TABLETS

NIGHT

**ASPIRIN (NSAID)** / Pain Reliever-Fever Reducer


Dextromethorphan HBr / Cough Suppressant

Doxylamine Succinate / Antihistamine

Phenylephrine Bitartrate / Nasal Decongestant

- Nasal Congestion
- Headache & Body Ache
- Cough
- Runny nose
- Sore Throat

8 EFFERVESCENT TABLETS



<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>● redness or swelling is present</li> <li>● new symptoms occur</li> <li>● ringing in the ears or loss of hearing occurs</li> <li>● cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> <li>● nervousness, dizziness, or sleeplessness occurs</li> </ul> <p><small>If experienced at home, discontinue use &amp; consult a health</small></p>	<p><b>Drug Facts (continued)</b></p> <p>operating machinery</p> <ul style="list-style-type: none"> <li>● <b>irritability may occur, especially in children</b></li> </ul> <p>Stop use and ask a doctor if:</p> <ul style="list-style-type: none"> <li>● you experience any of the following signs of stomach bleeding</li> <li>● feel faint</li> <li>● vomit blood</li> <li>● have bloody or black stools</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>● minor aches and pains</li> <li>● headache</li> <li>● nasal and sinus congestion</li> <li>● temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. <small>Labels on this product are for information in the back of the box.</small></p>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>● minor aches and pains</li> <li>● headache</li> <li>● nasal congestion and pressure</li> <li>● cough</li> <li>● sore throat</li> <li>● temporarily reduces fever</li> </ul> <p><b>Warnings</b></p> <p><b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. <small>Labels on this product are for information in the back of the box.</small></p>
---	--	--	---



<p><b>It is important to use aspirin carefully, even in children.</b></p> <p><b>Professional before use.</b> It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.</li> <li>children under 12 years: do not use</li> </ul>	<ul style="list-style-type: none"> <li>have stomach pain that does not get better</li> <li>pain, cough, or nasal congestion gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.</li> <li>nervousness, dizziness, or sleeplessness occurs</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it</p>	<p><b>STRICTLY WARNING: THIS PRODUCT IS A NONSTEROIDAL ANTI-INFLAMMATORY DRUG (NSAID).</b></p> <p>with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>swelling (edema)</li> <li>asthma (wheezing)</li> <li>stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or blood thinning drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> </ul>	<p><b>STRICTLY WARNING: THIS PRODUCT IS A NONSTEROIDAL ANTI-INFLAMMATORY DRUG (NSAID).</b></p> <p>If-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p><b>Allergy alert:</b> Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> <li>hives</li> <li>swelling (edema)</li> <li>asthma (wheezing)</li> <li>stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older</li> <li>have had stomach ulcers or bleeding problems</li> <li>take a blood thinning (anticoagulant) or blood thinning drug</li> <li>take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</li> </ul>
--	--	--	--

<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Some throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you are allergic to aspirin or any other pain reliever/fever reducer</li> <li>if you are now taking a prescription nonsteroidal anti-inflammatory drug (NSAID) (contains drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the NSAID drug. If you do not know if your prescription drug contains an NSAID, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients</li> <li>in children under 12 years of age</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>stomach bleeding warning applies to you</li> <li>you have a history of stomach problems, such as heartburn</li> <li>you have high blood pressure, heart</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p>nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)</p> <ul style="list-style-type: none"> <li>have 3 or more alcoholic drinks every day while using this product</li> <li>take more or for a longer time than directed</li> </ul> <p><b>Some throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use to sedate children.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you are allergic to aspirin or any other pain reliever/fever reducer</li> <li>if you are now taking a prescription nonsteroidal anti-inflammatory drug (NSAID) (contains drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the NSAID drug. If you do not know if your prescription drug contains an NSAID, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients</li> <li>in children under 12 years of age</li> </ul> <p><b>Ask a doctor before use if</b></p> <ul style="list-style-type: none"> <li>stomach bleeding warning applies to you</li> <li>you have a history of stomach problems,</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p>disease, liver cirrhosis, or kidney disease</p> <ul style="list-style-type: none"> <li>you are taking a diuretic</li> <li>you have</li> <li>asthma</li> <li>thyroid disease</li> <li>diabetes</li> <li>cough with excessive phlegm (mucus)</li> <li>difficulty in urination due to enlargement of the prostate gland</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>a sodium-restricted diet</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>taking a prescription drug for</li> <li>gout</li> <li>diabetes</li> <li>arthritis</li> </ul> <p><b>When using this product do not exceed recommended dosage</b></p> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>an allergic reaction occurs. Seek medical help right away.</li> <li>you experience any of the following signs of stomach bleeding</li> <li>feel faint</li> <li>feel hot</li> <li>feel blood</li> <li>have bloody or black stools</li> <li>have stomach pain that does not get better</li> <li>stomach bleeding warning applies to you</li> <li>you have a history of stomach problems,</li> </ul>	<p><b>Drug Facts (continued)</b></p> <p>such as heartburn</p> <ul style="list-style-type: none"> <li>you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</li> <li>you are taking a diuretic</li> <li>you have</li> <li>asthma</li> <li>thyroid disease</li> <li>diabetes</li> <li>glaucoma</li> <li>cough with excessive phlegm (mucus)</li> <li>a breathing problem such as emphysema or chronic bronchitis</li> <li>difficulty in urination due to enlargement of the prostate gland</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>a sodium-restricted diet</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>taking a prescription drug for</li> <li>gout</li> <li>diabetes</li> <li>arthritis</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>do not exceed recommended dosage</li> <li>may cause medical drowsiness</li> <li>avoid alcoholic drinks</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> <li>be careful when driving a motor vehicle or</li> </ul>
--	--	--	---

<p><b>Drug Facts (continued)</b></p> <p><b>Other Information</b></p> <ul style="list-style-type: none"> <li>each tablet contains sodium 416 mg</li> <li>Phenylephrine: Contains Phenylephrine 5 mg Per Tablet</li> <li>store at room temperature. Avoid excessive heat.</li> </ul> <p><b>Inactive Ingredients</b> acetylsalicylic acid, anhydrous citric acid, aspartame, calcium silicate, dextromethorphan, doxylamine, sodium, flavors, menthol, povidone, sodium benzoate, sodium bicarbonate</p> <p><b>Questions or comments?</b> 1-800-985-6085 (Mon-Fri 9AM-5PM EST)</p>	<p><b>Drug Facts (continued)</b></p> <p>may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water at bedtime (may be taken every 4 to 6 hours). Do not exceed 8 tablets in 24 hours or as directed by a doctor.</li> <li>children under 12 years: do not use</li> </ul> <p><b>Other Information</b></p> <ul style="list-style-type: none"> <li>each tablet contains: sodium 476 mg</li> <li>Phenylephrine: Contains Phenylephrine 5.6 mg Per Tablet</li> <li>store at room temperature. Avoid excessive heat.</li> </ul> <p><b>Inactive Ingredients</b> acetylsalicylic acid, anhydrous citric acid, aspartame, calcium silicate, dextromethorphan, doxylamine, sodium, flavors (natural &amp; artificial), menthol, povidone, sodium benzoate, sodium bicarbonate</p> <p><b>Questions?</b> 1-800-985-6085</p>
---	--

# ALKA-SELTZER PLUS COLD DAY AND NIGHT

aspirin, dextromethorphan hydrobromide, phenylephrine bitartrate, doxylamine succinate kit

## Product Information

Product Type		HUMAN OTC DRUG	Item Code (Source)		NDC:0280-1550
Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0280-1550-20	1 in 1 CARTON; Type 0: Not a Combination Product	09/14/2018		
Quantity of Parts					
Part #	Package Quantity		Total Product Quantity		
Part 1	6 POUCH		12		
Part 2	4 POUCH		8		
Part 1 of 2					
ALKA-SELTZER PLUS COLD DAY					
aspirin, dextromethorphan hydrobromide, phenylephrine bitartrate tablet, effervescent					
Product Information					
Route of Administration		ORAL			
Active Ingredient/Active Moiety					
Ingredient Name			Basis of Strength	Strength	
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)			PHENYLEPHRINE BITARTRATE	7.8 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)			ASPIRIN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)			DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
Inactive Ingredients					
Ingredient Name				Strength	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)					
CALCIUM SILICATE (UNII: S4255P4G5M)					
SODIUM BENZOATE (UNII: OJ245FE5EU)					
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)					
DOCUSATE SODIUM (UNII: F05Q2T2JA0)					
MANNITOL (UNII: 3OWL53L36A)					
POVIDONE (UNII: FZ989GH94E)					
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)					
ASPARTAME (UNII: Z0H242BBR1)					
DIMETHICONE (UNII: 92RU3N3Y1O)					
Product Characteristics					

Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	CITRUS	Imprint Code	ASP;DAY
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		6 in 1 CARTON		
1		2 in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/14/2018	

### Part 2 of 2

### ALKA-SELTZER PLUS COLD NIGHT

aspirin, dextromethorphan hydrobromide, phenylephrine bitartrate, doxylamine succinate tablet, effervescent

### Product Information

Route of Administration	ORAL
-------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
PHENYLEPHRINE BITARTRATE (UNII: 27O3Q5ML57) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

### Inactive Ingredients

Ingredient Name	Strength
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CALCIUM SILICATE (UNII: S4255P4G5M)	
ASPARTAME (UNII: Z0H242BBR1)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
MANNITOL (UNII: 3OWL53L36A)	

DIMETHICONE (UNII: 92RU3N3Y1O)				
<b>Product Characteristics</b>				
<b>Color</b>	white	<b>Score</b>	no score	
<b>Shape</b>	ROUND	<b>Size</b>	25mm	
<b>Flavor</b>	LEMON	<b>Imprint Code</b>	ASP;NT	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1		4 in 1 CARTON		
1		2 in 1 POUCH; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341		09/14/2018	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part341		09/14/2018	

**Labeler** - Bayer HealthCare LLC. (112117283)